

## ***Puccinia gladioli* on *Gladiolus***

*Puccinia gladioli* is a heteroecious rust of quarantine significance for the U.S. It is known from Europe and Asia. Although the aecial stage on *Valerianella* is reported from the western U.S., no telia have been found on *Gladiolus* in the U.S.

***Puccinia gladioli*** Castagne, Obs. Pl. Acotyl. 2:17 1843.

On *Valerianella*:

**Spermogonia** generally hypophyllous, yellow, eventually blackish, globose, with periphyses.

**Aecia** generally hypophyllous, scattered, occasionally epiphyllous and isolated, cylindric or cupulate, 250-300 µm diam., peridium margin lacerate, peridial cells rhomboid, 20-34 x 17-24 µm, external wall transversally striate, 3-6 µm thick, internal wall verrucose, 2-3.5 µm thick; aeciospores subglobose, angular globose to ellipsoid, 17-24 x 13-18 µm, wall 1.5-3 µm thick, minutely verrucose.

On *Gladiolus*:

**Uredinia** unknown

**Telia** amphigenous, on reddish spots, sometimes limited by veins, minute, rounded, densely crowded or confluent, forming a crust up to 1 cm long, often covering much of the leaf surface, compact, chestnut brown to black; paraphyses cylindric to slightly clavate, brown, up to 80 µm long; teliospores ellipsoid to clavate, apex round to acute, slightly constricted at septum, gradually narrowing below septum, 36-60 x 16-27 µm, wall smooth, pale brown, 2-3 µm thick, up to 10 µm at the apex, sometimes mesospores present, 24-40 x 12-17 µm; pedicel hyaline, persistent, 10-60 µm long.

**Hosts:** Mainly on species of *Gladiolus* and *Valerianella*

**Geographic distribution:** Reported on *Gladiolus* from Europe and Asia. Although telia of *Puccinia gladioli* on *Gladiolus* have not been reported from the U.S., the aecial stage (*Aecidium valerianellae*) on *Valerianella* has been reported from the western U.S., as well Europe and Asia.


The heteroecious condition of this rust was proven by d'Oliviera (1949) who inoculated aeciospores of *Aecidium valerianellae* from *Valerianella* onto *Gladiolus* and observed production of telia typical of *P. gladioli* after 10 days (Wilson and Henderson 1966).

The only other *Puccinia* reported on *Gladiolus* is *Puccinia mccleanii* Doidge (1941) from South Africa. That species produces paraphysate telia on *Gladiolus* whereas *P. gladioli* produces telia with paraphyses. Teliospores of *P. mccleanii* are thinner and paler than those of *P. gladioli*, which produces teliospores with cell walls usually thickened at the apex, up to 10 µm.

### **References:**

- d'Oliviera, B.** 1949. Life cycle of *Puccinia gladioli* Cast. Nature 164: 239.  
**Doidge, E.M.** 1941. South African rust fungi IV. Bothalia 4: 229-236.

Aecia BPI 153567



Aecia BPI 153566




Aecia BPI 153566



Aeciospores BPI 153570




Aeciospores BPI 153570



Aeciospores BPI 153570




Telia BPI 068188




Teliospores BPI 068187



Teliospores BPI 068188



Telia BPI 068188



Teliospores BPI 068187

